

Other Squashing Deformation and $\mathcal{N} = 3$ Superconformal Chern-Simons Gauge Theory

Changhyun Ahn

Department of Physics, Kyungpook National University, Taegu 702-701, Korea

ahn@knu.ac.kr

Abstract

We consider one of the well-known solutions in eleven-dimensional supergravity where the seven-dimensional Einstein space is given by a $SO(3)$ -bundle over the $\mathbf{CP}^2$. By reexamining the AdS_4 supergravity scalar potential, the holographic renormalization group flow from $\mathcal{N} = (0, 1)$ $SU(3) \times SU(2)$ -invariant UV fixed point to $\mathcal{N} = (3, 0)$ $SU(3) \times SU(2)$ -invariant IR fixed point is reinterpreted. A dual operator in three-dimensional superconformal Chern-Simons matter theories corresponding to this RG flow is described.

1 Introduction

The $\mathcal{N} = 6$ superconformal Chern-Simons matter theory with gauge group $U(N) \times U(N)$ at level k and with two hypermultiplets in the bifundamental representations is found in [1]. This gauge theory is described as the low energy limit of N M2-branes probing $\mathbf{C}^4/\mathbf{Z}_k$ singularity. At large N -limit, this theory is dual to the eleven-dimensional M-theory on $AdS_4 \times \mathbf{S}^7/\mathbf{Z}_k$ where the seven-sphere metric is realized as an $\mathbf{S}^1$ -fibration over $\mathbf{CP}^3$ [2]. One of the main observations in [1] is to look at the special case of $\mathcal{N} = 3$ superconformal Chern-Simons matter theory with above particular gauge group, matter contents, and particular choice of Chern-Simons levels of two gauge groups. Then naive $SU(2)$ flavor symmetry appearing in the hypermultiplets is enhanced to $SU(2) \times SU(2)$ symmetry that occurs in the whole action of the theory and there exists a $SU(2)_R$ symmetry coming from the original $\mathcal{N} = 3$ superconformal symmetry. It turns out the full theory has $\mathcal{N} = 6$ superconformal symmetry and the full scalar potential is invariant under $SU(4)_R$ coming from this enhanced $\mathcal{N} = 6$ superconformal symmetry.

The simplest spontaneous compactification of the eleven-dimensional supergravity is the Freund-Rubin [3] compactification to a product of AdS_4 spacetime and an arbitrary compact seven-dimensional Einstein manifold $\mathbf{X}^7$ of positive scalar curvature. The standard Einstein metric of the round seven-sphere $\mathbf{S}^7$ yields a vacuum with $SO(8)$ gauge symmetry and $\mathcal{N} = 8$ supersymmetry. There exists a second squashed Einstein metric [4, 5] yielding a vacuum with $SO(5) \times SU(2)$ gauge symmetry and $\mathcal{N} = (1, 0)$ supersymmetry [6, 7]. As suggested in [8, 9], in [10], it was shown that the well-known spontaneous (super)symmetry breaking deformation from round $\mathbf{S}^7$ to squashed one is mapped to a renormalization group(RG) flow from $\mathcal{N} = (1, 0)$ $SO(5) \times SU(2)$ -invariant fixed point in the UV to $\mathcal{N} = 8$ $SO(8)$ -invariant fixed point in the IR. In particular, the squashing deformation corresponds to an irrelevant operator at the UV superconformal fixed point and a relevant operator at the IR (super)conformal fixed point respectively. Moreover the RG flow is described geometrically by a static domain wall which interpolates the two asymptotically AdS_4 spacetimes with round and squashed $\mathbf{S}^7$'s. For the different type of compactifications where the internal space has nonzero four-form field strength, see also [11, 12].

One could ask [13] what happens when we perform $\mathbf{Z}_k$ -quotient [1] along the above whole RG flow [10]? Starting from the general, one parameter-family, metric for $\mathbf{CP}^3$ inside of seven-sphere and its seven-dimensional uplift metric on an $\mathbf{S}^1$ -bundle over this $\mathbf{CP}^3$, the full eleven-dimensional metric with appropriate warp factors was constructed. By analyzing the AdS_4 scalar potential, the holographic supersymmetric(or nonsupersymmetric) RG flow from $\mathcal{N} =$

$(1, 0)$ $SO(5) \times U(1)$ -invariant UV fixed point to $\mathcal{N} = (6, 0)$ $SU(4)_R \times U(1)$ -invariant IR fixed point was described in [13]. Each symmetry group is the subset of previous ones respectively. That is, $SO(5) \times U(1)$ is contained in $SO(5) \times SU(2)$ and $SU(4)_R \times U(1)$ is contained in $SO(8)$. The squashing deformation corresponds to the singlet of $\mathbf{20}'$ of $SU(4)_R$ and it is given by the quartic term for the matter fields transforming as fundamental representation under the $SU(4)_R$. The dual Chern-Simons matter theory at the $\mathcal{N} = (1, 0)$ $SO(5) \times U(1)$ -invariant UV fixed point is constructed in [14].

Now it is natural to ask that are there any other examples where some squashing deformation in $\mathbf{X}^7$ might provide a similar RG flow and one can think of some dual operator in three-dimensional boundary conformal field theory? Yes, the manifold $\mathbf{X}^7 = N_I^{0,1,0}$ has been studied originally by Castellani and Romans [15] who identified this manifold as a particular coset manifold which is specified by three integers. We consider the particular case where $p = 0, q = 1$ and $r = 0$ in $N_I^{p,q,r}$ manifold. There exists $\mathcal{N} = (3, 0)$ supersymmetry with $SU(3) \times SU(2)$ gauge symmetry or $\mathcal{N} = (1, 0)$ supersymmetry with $SU(3) \times U(1)$ gauge symmetry (See also [16]). Moreover, Page and Pope [17] have completed the coset manifold construction by showing existence of another family of Einstein manifold, $N_{II}^{0,1,0}$, which can be obtained from geometric squashing of the $N_I^{0,1,0}$, retains the same gauge group $SU(3) \times SU(2)$ but instead preserves $\mathcal{N} = (0, 1)$ supersymmetry. As in the case of seven-sphere $\mathbf{S}^7$, the scalar field corresponding to the squashing deformation acquires a nonzero vacuum expectation value leading to (super)-Higgs mechanism. With left-orientation the squashing interpolates between a $\mathcal{N} = 3$ supersymmetric vacuum and another with $\mathcal{N} = 0$ supersymmetry. With right-orientation it interpolates between a nonsupersymmetric vacuum and a supersymmetry restored one with $\mathcal{N} = 1$ supersymmetry [18].

On the other hand, in [19], the corresponding $\mathcal{N} = 3$ dual gauge theory has gauge group $SU(N) \times SU(N)$ with “three” hypermultiplets transforming as a “triplet” under the $SU(3)$ flavor symmetry which is nothing but one of the global symmetries for $N_I^{0,1,0}$ manifold¹. In terms of $\mathcal{N} = 2$ superfields, these hypermultiplets can be reorganized as two sets of chiral superfields. For the color representation, one of these transforms as $(\mathbf{N}, \overline{\mathbf{N}})$ and the other transforms as $(\overline{\mathbf{N}}, \mathbf{N})$. Furthermore, these two superfields transform as a doublet of $SU(2)_R$ which is the $SU(2)$ factor in the remaining global symmetry of $N_I^{0,1,0}$ manifold. As we mentioned before, $SU(2)_R$ corresponds to the $\mathcal{N} = 3$ superconformal symmetry. After integrating out two adjoint fields of the theory, the effective quartic superpotential can be

¹We emphasize that this theory is based on Yang-Mills plus Chern-Simons theory with chiral multiplets, contrary to [1] in which the theory is described as Chern-Simons theory with chiral multiplets. For example, the kinetic terms in [19] contain those for the vector multiplet as well as the Chern-Simons term and those for the chiral multiplets. However, in [1], there is no kinetic term for any of the fields in the vector multiplet.

obtained. The coefficient is determined by the $\mathcal{N} = 3$ supersymmetry but breaks $\mathcal{N} = 4$ supersymmetry. For the general discussion on $\mathcal{N} = 3$ superconformal Chern-Simons matter theory, see [20]. The complete $\mathcal{N} = 3$ Kaluza-Klein spectrum is found in [21] and its $OSp(3|4)$ multiplet structure is further explained in [22].

Later, Billo, Fabbri, Fre, Merlatti and Zaffaroni [23](See also [24]) have constructed an $\mathcal{N} = 3$ long massive spin $3/2$ multiplet with conformal dimension 3 from the massless $\mathcal{N} = 3$ graviton multiplet. These two are connected to “shadow” relation: fields of different type, spin and mass are linked by a relation which determines the mass of the one as a function of the other.

In this paper, we will be studying the known example of Kaluza-Klein supergravity vacua and reinterpret it in terms of three-dimensional (super)conformal field theories and associated RG flows. We will be exploring the Freund-Rubin type spontaneous compactification on $AdS_4 \times \mathbf{X}^7$. For M2-branes on an eight-dimensional manifold, the near-horizon geometry $\mathbf{X}^7$ is expected to change as the M2-branes are placed at or away a conical singularity of the manifold [25, 26]. More specifically, we will consider $\mathbf{X}^7$ being 3-Sasaki holonomy manifold, describing near-horizon geometry of M2-branes at relevant conical singularities.

When the work of [18] was completed at that time, it was not possible to analyze the gauge theory description because the Kaluza-Klein spectrum of eleven-dimensional supergravity was not complete. Later, in [23], they have found more mass spectrum in the eleven-dimensional supergravity side that includes the harmonics of the Lichnerowicz scalar with conformal dimension 4. Then one can identify the corresponding fluctuation spectrum for the scalar fields around $\mathcal{N} = 3$ fixed point.

The aim of this paper is to 1) recapitulate the effective scalar potential described in [18] with only breathing mode and squashing mode, and 2) analyze more both the mass spectrum in the eleven-dimensional supergravity and the corresponding Chern-Simons gauge theory operator which gives rise to the squashing deformation, by analyzing the results of [23].

In section 2, we describe the seven-dimensional Einstein space($N_I^{0,1,0}$) and its squashed version($N_{II}^{0,1,0}$) compactification vacua in eleven-dimensional supergravity. The effective four-dimensional scalar potential looks similar to the one for seven-sphere and the two critical points have nonzero scalar fields. However, the ratio of the squashing parameter at these two critical values(which is equal to $1/5$) is the same as the one in seven-sphere case.

In section 3, the squashing deformation of each vacua is described by an irrelevant operator at the $\mathcal{N} = (3, 0)$ conformal fixed point and a relevant operator at the $\mathcal{N} = (0, 1)$ conformal fixed points. The RG flow is described in AdS_4 supergravity by a static domain wall interpolating between these two vacua. We identify the corresponding operator in the

boundary conformal field theory in three dimensions by looking at the observations of [23]².

2 Two seven-dimensional Einstein spaces

A generic eleven-dimensional metric interpolating between two seven-dimensional Einstein spaces with an arbitrary four-dimensional spacetime metric maybe written as

$$\begin{aligned} \frac{\overline{ds}^2}{R^2} = & e^{-7u} g_{\alpha\beta} dx^\alpha dx^\beta + e^{2u+3v} \left[d\mu^2 + \frac{1}{4} \sin^2 \mu (\sigma_1^2 + \sigma_2^2 + \cos^2 \mu \sigma_3^2) \right] \\ & + e^{2u-4v} \left[(\Sigma_1 - \cos \mu \sigma_1)^2 + (\Sigma_2 - \cos \mu \sigma_2)^2 + \left(\Sigma_3 - \frac{1}{2}(1 + \cos^2 \mu) \sigma_3 \right)^2 \right], \end{aligned} \quad (2.1)$$

where the three real left-invariant one-forms satisfy the $SU(2)$ algebra $d\sigma_i = -\frac{1}{2}\epsilon_{ijk}\sigma_j \wedge \sigma_k$ and those on the manifold $SO(3)$ which is necessary for the regularity of the metric have $d\Sigma_i = -\frac{1}{2}\epsilon_{ijk}\Sigma_j \wedge \Sigma_k$ ³. The coordinate μ and $SU(2)$ one-forms σ_i are the same as $\mathbf{CP}^2$ metric and corresponding isometry is characterized by $SU(3)$. Then the seven-dimensional $N^{0,1,0}$ space is a nontrivial $SO(3)$ bundle over $\mathbf{CP}^2$ and the isometry group for this space is $SU(3) \times SU(2)$ [17]. The scalar fields $u(x)$ for the breathing mode and $v(x)$ for squashing mode depend on the four-dimensional spacetime. Moreover, the squashing is parametrized by [34]

$$\lambda^2 \equiv e^{-7v}. \quad (2.2)$$

The parameter R measures the overall radius of curvature. The gauge fields are given by $A_1 = \cos \mu \sigma_1$, $A_2 = \cos \mu \sigma_2$ and $A_3 = \frac{1}{2}(1 + \cos^2 \mu) \sigma_3$.

Spontaneous compactification of M-theory to $AdS_4 \times \mathbf{X}^7$ is obtained from near-horizon geometry of N coincident M2-branes and the nonvanishing flux of four-form field strength of the Freund-Rubin [3] is given by

$$\overline{F}_{\alpha\beta\gamma\delta} = Q e^{-7u} \overline{\epsilon}_{\alpha\beta\gamma\delta} = Q e^{-21u} \epsilon_{\alpha\beta\gamma\delta}. \quad (2.3)$$

Here the Page charge [34, 32] defined by $Q \equiv \pi^{-4} \int_{\mathbf{X}^7} (*\overline{F} + \overline{C} \wedge \overline{F})$ is related to the total number of M2-branes through $Q = 96\pi^2 N \ell_p^6$. Then the eleven-dimensional Einstein equation

²Recently, the $\mathcal{N} = 3$ superconformal Chern-Simons quiver theories are constructed in [27] but these theories do not contain the seven-dimensional Einstein manifold we are considering here. See also other relevant paper [28]. For the earlier studies on $\mathcal{N} = 3$ superconformal Chern-Simons theories, there are also some works in [29, 30, 31].

³ They are given by $\sigma_1 = \cos \psi d\theta + \sin \psi \sin \theta d\phi$, $\sigma_2 = -\sin \psi d\theta + \cos \psi \sin \theta d\phi$ and $\sigma_3 = d\psi + \cos \theta d\phi$ and similarly $SO(3)$ one-forms are given by $\Sigma_1 = \cos \gamma d\alpha + \sin \gamma \sin \alpha d\beta$, $\Sigma_2 = -\sin \gamma d\alpha + \cos \gamma \sin \alpha d\beta$, and $\Sigma_3 = d\gamma + \cos \alpha d\beta$ [32, 33].

with (2.3) provides the following Ricci tensor components [34, 13]

$$\overline{R}_\beta^\alpha = -\frac{4}{3}Q^2 e^{-14u} \delta_\beta^\alpha, \quad \overline{R}_b^a = \frac{2}{3}Q^2 e^{-14u} \delta_b^a, \quad \overline{R}_b^\alpha = \overline{R}_\beta^a = 0. \quad (2.4)$$

On the other hand, the Ricci tensor components can be obtained from the following orthonormal basis which can be read off from the eleven-dimensional metric (2.1)

$$\begin{aligned} e^1 &= e^{-\frac{7}{2}u} \sqrt{g_{11}(x)} dx^1, & e^2 &= e^{-\frac{7}{2}u} \sqrt{g_{22}(x)} dx^2, \\ e^3 &= e^{-\frac{7}{2}u} \sqrt{g_{33}(x)} dx^3, & e^4 &= e^{-\frac{7}{2}u} \sqrt{g_{44}(x)} dx^4, \\ e^5 &= e^{u+\frac{3}{2}v} d\mu, & e^6 &= \frac{1}{2} e^{u+\frac{3}{2}v} \sin \mu \sigma_1, \\ e^7 &= \frac{1}{2} e^{u+\frac{3}{2}v} \sin \mu \sigma_2, & e^8 &= \frac{1}{2} e^{u+\frac{3}{2}v} \sin \mu \cos \mu \sigma_3, \\ e^9 &= e^{u-2v} (\Sigma_1 - \cos \mu \sigma_1), & e^{10} &= e^{u-2v} \sin \mu (\Sigma_2 - \cos \mu \sigma_2), \\ e^{11} &= e^{u-2v} \left[\Sigma_3 - \frac{1}{2} (1 + \cos^2 \mu) \sigma_3 \right]. \end{aligned} \quad (2.5)$$

The results for Ricci tensor in the basis of (2.5) are summarized by

$$\begin{aligned} \overline{R}_\beta^\alpha &= e^{7u} \left(R_\beta^\alpha + \frac{7}{2} \delta_\beta^\alpha u_{;\gamma}^{\gamma} - \frac{63}{2} u^{;\alpha} u_{;\beta} - 21 v^{;\alpha} v_{;\beta} \right), \\ \overline{R}_5^5 &= 6e^{-2u-3v} - 6e^{-2u-10v} - e^{7u} \left(u_{;\alpha}^{\alpha} + \frac{3}{2} v_{;\alpha}^{\alpha} \right) = \overline{R}_6^6 = \overline{R}_7^7 = \overline{R}_8^8, \\ \overline{R}_9^9 &= \frac{1}{2} e^{-2u+4v} + 4e^{-2u-10v} - e^{7u} (u_{;\alpha}^{\alpha} - 2v_{;\alpha}^{\alpha}) = \overline{R}_{10}^{10} = \overline{R}_{11}^{11}. \end{aligned} \quad (2.6)$$

Note that the only first terms in $\overline{R}_5^5$ and $\overline{R}_9^9$ are different from the one for squashed $\mathbf{S}^7$ space [34, 13]⁴.

Substituting the last two relations in (2.6) into (2.4) implies the field equations for the breathing mode $u(x)$ and the squashing mode $v(x)$ as follows:

$$\begin{aligned} u_{;\alpha}^{\alpha} &= \frac{3}{14} e^{-9u+4v} + \frac{24}{7} e^{-9u-3v} - \frac{12}{7} e^{-9u-10v} - \frac{2}{3} Q^2 e^{-21u}, \\ v_{;\alpha}^{\alpha} &= -e^{-9u+4v} + 12e^{-9u-3v} - 20e^{-9u-10v}. \end{aligned} \quad (2.7)$$

Note that the only first two terms in $u_{;\alpha}^{\alpha}$ and $v_{;\alpha}^{\alpha}$ are different from the one for squashed $\mathbf{S}^7$ space [34, 13]. The vanishing of the right hand side of second equation of (2.7) implies that either $v = v_1 = \frac{1}{7} \ln 2$ or $v = v_2 = \frac{1}{7} \ln 10$. Furthermore, substituting the field equation

⁴ When the scalar fields $u(x)$ and $v(x)$ are constant, then $\overline{R}_5^5 = 6 - 6e^{-7v} = 6 - 6\lambda^2 = \overline{R}_6^6 = \overline{R}_7^7 = \overline{R}_8^8$ and $\overline{R}_9^9 = 4e^{-7v} + \frac{1}{2}e^{7v} = 4\lambda^2 + \frac{1}{2\lambda^2} = \overline{R}_{10}^{10} = \overline{R}_{11}^{11}$ where (2.2) is used. Then $\lambda^2 = \frac{1}{2}$ corresponds to the Einstein metric in [15] and $\lambda^2 = \frac{1}{10}$ corresponds to the squashed Einstein metric [17].

for $u(x)$ in (2.7) into the first equation of (2.6) together with (2.4) implies the following four-dimensional Ricci tensor

$$R_{\beta}^{\alpha} = \frac{63}{2} u^{;\alpha} u_{;\beta} + 21 v^{;\alpha} v_{;\beta} + \delta_{\beta}^{\alpha} e^{-9u} \left(-\frac{3}{4} e^{4v} - 12 e^{-3v} + 6 e^{-10v} + Q^2 e^{-12u} \right). \quad (2.8)$$

Now the field equations (2.7) and (2.8) are equivalent to the Euler-Lagrange equations for the following effective Lagrangian

$$\mathcal{L} = \sqrt{-g} \left[R - \frac{63}{2} (\partial u)^2 - 21 (\partial v)^2 - V(u, v) \right] \quad (2.9)$$

with the scalar potential ⁵

$$V(u, v) = e^{-9u} \left(-\frac{3}{2} e^{4v} - 24 e^{-3v} + 12 e^{-10v} + 2 Q^2 e^{-12u} \right). \quad (2.10)$$

Note that the first two terms in (2.10) are different from the one for squashed $\mathbf{S}^7$ space [34, 10, 13].

One analyzes two vacua of this scalar potential as follows:

$$\begin{aligned} N_{\text{I}}^{0,1,0} \quad : \quad u = u_1 &= \frac{1}{12} \ln \left(\frac{2^{\frac{10}{7}}}{3^2} Q^2 \right), \quad v = v_1 = \frac{1}{7} \ln 2, \quad (\lambda^2 = \frac{1}{2}), \\ \Lambda_1 &= -9 \left| \frac{2Q^3}{3} \right|^{-\frac{1}{2}}, \end{aligned}$$

and

$$\begin{aligned} N_{\text{II}}^{0,1,0} \quad : \quad u = u_2 &= \frac{1}{12} \ln \left(\frac{10^{\frac{10}{7}}}{3^4} Q^2 \right), \quad v = v_2 = \frac{1}{7} \ln 10, \quad \left(\lambda^2 = \frac{1}{10} \right), \\ \Lambda_2 &= -3^6 \cdot 5^{-2} |10Q^3|^{-\frac{1}{2}}. \end{aligned}$$

The two supergravity solutions are classically stable under the changes of the size and squashing parameter of seven-dimensional space [36]. The $N_{\text{I}}^{0,1,0}$ is a saddle point, corresponds to a minimum along the v -direction and is invariant under the $SU(3) \times SU(2)$ isometry group while $N_{\text{II}}^{0,1,0}$ is a maximum and is invariant under the same $SU(3) \times SU(2)$ isometry group. The left-handed seven-dimensional space $N_{\text{I,L}}^{0,1,0}$ gives rise to a theory with $\mathcal{N} = 3$ supersymmetry while the right-handed seven-dimensional space $N_{\text{I,R}}^{0,1,0}$ gives rise to a theory with no supersymmetry ($\mathcal{N} = 0$) [17].

Moreover, the left-handed squashed seven-dimensional space $N_{\text{II,L}}^{0,1,0}$ gives rise to a theory with no supersymmetry while the right-handed squashed seven-dimensional space $N_{\text{II,R}}^{0,1,0}$ gives

⁵This form for the scalar potential was observed also previously in [35].

rise to a theory with $\mathcal{N} = 1$ supersymmetry. That is, with the choice of left-handed orientation of $N_L^{0,1,0}$, one regards the $\lambda^2 = \frac{1}{2}$ metric as giving the unbroken vacuum state with $\mathcal{N} = 3$ supersymmetry which can be broken spontaneously to $\lambda^2 = \frac{1}{10}$ metric yielding $\mathcal{N} = 0$ supersymmetry. On the other hand, with the choice of opposite orientation of $N_R^{0,1,0}$ the $\lambda^2 = \frac{1}{10}$ metric provides a vacuum state with $\mathcal{N} = 1$ supersymmetry which can be broken to the $\lambda^2 = \frac{1}{2}$ metric with $\mathcal{N} = 0$ supersymmetry.

Therefore, for the squashing with left-handed orientation, the RG flow interpolates between the boundary conformal field theories with $\mathcal{N} = 3$ and $\mathcal{N} = 0$ supersymmetry while for the squashing with right-handed orientation, the RG flow interpolates between conformal field theories with $\mathcal{N} = 0$ and $\mathcal{N} = 1$ ⁶.

3 Three-dimensional (super)conformal field theories

Using the previous results on the Kaluza-Klein spectrum under squashing deformations, an operator giving rise to a RG flow associated with the supersymmetry breaking will be identified and it turns out that the operator is relevant at the $N_{\text{II}}^{0,1,0}$ fixed point and irrelevant at the $N_{\text{I}}^{0,1,0}$ fixed point with same $SU(3) \times SU(2)$ symmetry groups.

- $SU(3) \times SU(2)$ -invariant conformal fixed point

Let us consider the harmonic fluctuations of spacetime metric and $u(x)$ and $v(x)$ scalar fields around $AdS_4 \times N_{\text{I}}^{0,1,0}$. Following [34, 10, 13], it is more convenient to rewrite (2.9) in terms of the unrescaled M-theory metric $\overline{g}_{\alpha\beta} = e^{-7u} g_{\alpha\beta}$ in (2.1):

$$\mathcal{L} = \sqrt{-\overline{g}} e^{7u} \left[\overline{R} - 2\overline{\Lambda}_1 - 105(\partial u)^2 - 21(\partial v)^2 - 2\overline{V}_1(u, v) \right], \quad (3.1)$$

where the scalar potential is written as

$$\overline{V}_1(u, v) = -\overline{\Lambda}_1 \left[1 - \frac{1}{4} e^{-2(u-u_1)} (e^{4(v-v_1)} + 8e^{-3(v-v_1)} - 2e^{-10(v-v_1)}) + \frac{3}{4} e^{-14(u-u_1)} \right]$$

⁶It is known that the $N_{\text{I}}^{0,1,0}$ can be written as an $U(1)$ -fibration over a Kahler-Einstein six-manifold that is an $\mathbf{S}^2$ -bundle over $\mathbf{CP}^2$ [37]. For the general squashed case $N^{0,1,0}$ including $N_{\text{II}}^{0,1,0}$, one can also get $U(1)$ -fibration over a six-manifold by reorganizing the last line of (2.1):

$$\begin{aligned} ds_7^2 = & 2\lambda^2 \left[-\frac{1}{2} d\psi - \frac{1}{2} \cos \mu \sin \theta (\cos \phi \sigma_1 - \sin \phi \sigma_2) + \frac{1}{4} \cos \theta (1 + \cos^2 \mu) \sigma_3 - \frac{1}{2} \cos \theta d\phi \right]^2 \\ & + \frac{\lambda^2}{2} \left([d\theta - \cos \mu (\sin \phi \sigma_1 + \cos \phi \sigma_2)]^2 + \sin^2 \theta \left[d\phi - \cos \mu \cot \theta (\cos \phi \sigma_1 - \sin \phi \sigma_2) - \frac{1}{2} (1 + \cos^2 \mu) \sigma_3 \right]^2 \right) \\ & + \frac{1}{2} \left[d\mu^2 + \frac{1}{4} \sin^2 \mu (\sigma_1^2 + \sigma_2^2 + \cos^2 \mu \sigma_3^2) \right], \end{aligned}$$

where we choose the same parametrization given in [37]. The Euler angles θ, ϕ and ψ correspond to the one-forms Σ_i . Of course, for the case of $\lambda^2 = \frac{1}{2}$, this metric leads to the one in [37].

in which the un-rescaled cosmological constant $\overline{\Lambda}_1 = e^{7u_1}\Lambda_1 = \frac{1}{2}e^{7u_1}V(u_1, v_1)$ is given by

$$\overline{\Lambda}_1 \equiv -3m_1^2 \frac{1}{\ell_p^2} = -3 \left(\frac{|Q|}{6} \right)^{-\frac{1}{3}} \frac{1}{\ell_p^2} \quad \text{where} \quad m_1 = \frac{1}{\overline{r}_{\text{IR}}}. \quad (3.2)$$

Here $\overline{r}_{\text{IR}}$ is related to N and Planck scale ℓ_p as $\overline{r}_{\text{IR}} = \ell_p \frac{1}{2}(32\pi^2 N)^{1/6}$.

By rescaling the scalar fields from the kinetic terms in the Lagrangian (3.1) as

$$\sqrt{210}u \equiv \overline{u}, \quad \sqrt{42}v \equiv \overline{v},$$

one obtains the fluctuation spectrum for $\overline{v}$ -field around the $N_I^{0,1,0}$ which takes a positive value:

$$M_{\overline{v}\overline{v}}^2(N_I^{0,1,0}) = \left(\frac{\partial^2}{\partial \overline{v}^2} 2\overline{V}_1 \right)_{\overline{u}=\overline{u}_1, \overline{v}=\overline{v}_1} = -\frac{4}{3}\overline{\Lambda}_1 \ell_p^2 = 4m_1^2 \quad (3.3)$$

where the relation (3.2) is used.

Recall that in the compactification of $AdS_4 \times \mathbf{S}^7$, the $\overline{v}$ -field represents the squashing of $\mathbf{S}^7$ and hence ought to correspond to **300** (that is the Young tableaux $\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$ of $SO(8)$ or the $SO(8)$ Dynkin label is given by $(\mathbf{0}, \mathbf{2}, \mathbf{0}, \mathbf{0})$): the lowest mode of the transverse, traceless symmetric tensor representation. The branching rule of the representation **300** in terms of $SO(7)$ Dynkin labels is given by [38, 39] $\mathbf{300}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \rightarrow \mathbf{27}(\begin{smallmatrix} \square & \square \end{smallmatrix}) \oplus \mathbf{105}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \oplus \mathbf{168}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})$ and the Lichnerowicz operator Δ_L in $SO(7)$ representation becomes **27** [32] which has a Dynkin label $(\mathbf{2}, \mathbf{0}, \mathbf{0})$ and its branching rule in terms of maximal subgroup $SU(4)$ Dynkin labels is

given by $\mathbf{27}(\begin{smallmatrix} \square & \square \end{smallmatrix}) \rightarrow \mathbf{1}(\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}) \oplus \mathbf{6}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \oplus \mathbf{20}'(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})$. Note that the **20'** in $SU(4)$ is represented by a traceless symmetric matrix and the squashing should correspond to nonzero expectation value for the **20'** of $SU(4)$. Then the branching rule of this **20'** in terms of $SU(3)$ which is a global symmetry for $N_I^{0,1,0}$ is given by

$$\mathbf{20}'(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \rightarrow \mathbf{6}(\begin{smallmatrix} \square & \square \end{smallmatrix}) \oplus \overline{\mathbf{6}}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \oplus \mathbf{8}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}).$$

Moreover, the $\overline{\mathbf{6}}$ in $SU(3)$ is represented by a traceless symmetric matrix and the squashing should eventually correspond to nonzero expectation value for the $\overline{\mathbf{6}}$ of $SU(3)$.

According to the nice observations of [23], the harmonics of Lichnerowicz scalars with conformal dimension $\Delta = 4$ has been obtained. The harmonics are eigenfunctions of the Lichnerowicz operator with eigenvalue $M_{\mathbf{200}} = 96m^2$ in their notation. Recall that the representation $(\mathbf{2}, \mathbf{0}, \mathbf{0})$ refers to $SO(7)$ representation **27**. With the help of [40], the value of $M_{\mathbf{200}}$ can be obtained from $M_{\frac{3}{2}, \frac{1}{2}, \frac{1}{2}}$ which is an eigenvalue for Rarita Schwinger operator and it turns out -16 . The representation $(\frac{3}{2}, \frac{1}{2}, \frac{1}{2})$ refers to $SO(7)$ representation **48**. The explicit

form is given by $M_{\mathbf{200}} = (M_{\frac{3}{2}\frac{1}{2}\frac{1}{2}} + 4)(M_{\frac{3}{2}\frac{1}{2}\frac{1}{2}} + 8)m^2$. By plugging $M_{\frac{3}{2}\frac{1}{2}\frac{1}{2}} = -16$, one gets $M_{\mathbf{200}} = 96m^2$. Note that m^2 is mass-squared parameter of a given AdS_4 spacetime and the mass of a scalar field ϕ in [40] is defined as $(\Delta_{AdS} + M_{\mathbf{200}} - 32m^2)\phi = 0$. Then $96m^2$ goes to $64m^2$ by subtracting $32m^2$ and by dividing out 16 further in order to compare with the usual normalization in AdS/CFT correspondence, one arrives that the correct result for the mass-squared is $4m^2$ which is the same normalization used in [18]. For example, see also the recent paper [41] for the normalization between the conformal dimension and the mass-squared term. Therefore, the mass-squared for the representation $\bar{\mathbf{6}}$ of $SU(3)$ is given by

$$M_{\bar{\mathbf{6}}}^2 = 4m_1^2,$$

which is exactly equal to (3.3). Note that the assignment of $SU(2)_R$ isospin J is related to the $SU(3)$ assignment and the R-symmetry group is the maximal $SU(2)$ subgroup of $SU(3)$. Under this embedding, the $SU(3)$ representations decompose into as follows:

$$\mathbf{1}, \bar{\mathbf{1}} \rightarrow J = 0, \quad \mathbf{3}, \bar{\mathbf{3}} \rightarrow J = 1, \quad \mathbf{6}, \bar{\mathbf{6}} \rightarrow J = 0 \oplus J = 2.$$

Or we have $\bar{\mathbf{6}}(\boxplus) = \bar{\mathbf{1}}(\boxminus) \oplus \bar{\mathbf{5}}(\boxminus)$ under the breaking of $SU(3)$ into the $SU(2)_R$.

One concludes that, in three-dimensional conformal field theory with $\mathcal{N} = 3$ supersymmetry, the $SU(3) \times SU(2)$ symmetric left-handed squashing should be an irrelevant perturbation of conformal dimension $\Delta = 4$. Note that this gives a nonsupersymmetric theory for the right-handed orientation for seven-dimensional Einstein manifold $N_{\text{I,R}}^{0,1,0}$.

- $SU(3) \times SU(2)$ -invariant conformal fixed point

Due to the skew-whipping, the theory will be either left-squashed $N_{\text{II,L}}^{0,1,0}$ with $\mathcal{N} = 0$ supersymmetry or right-squashed $N_{\text{II,R}}^{0,1,0}$ with $\mathcal{N} = 1$ supersymmetry. The isometry of the squashed seven-dimensional Einstein manifold is given by $SU(3) \times SU(2)$. In terms of the unrescaled M-theory metric, the Lagrangian (2.9) can be rewritten as

$$\mathcal{L} = \sqrt{-\bar{g}}e^{7u} [\bar{R} - 2\bar{\Lambda}_2 - 105(\partial u)^2 - 21(\partial v)^2 - 2\bar{V}_2(u, v)],$$

where the scalar potential is given by

$$\bar{V}_2(u, v) = -\bar{\Lambda}_2 \left[1 - \frac{1}{36}e^{-2(u-u_2)} (25e^{4(v-v_2)} + 40e^{-3(v-v_2)} - 2e^{-10(v-v_2)}) + \frac{3}{4}e^{-14(u-u_2)} \right],$$

and the un-rescaled cosmological constant $\bar{\Lambda}_2 = e^{7u_2}\Lambda_2 = \frac{1}{2}e^{7u_2}V(u_2, v_2)$ is given by

$$\bar{\Lambda}_2 \equiv -3m_2^2 \frac{1}{\ell_p^2} = -3 \left[3^{\frac{7}{3}} \cdot 5^{-\frac{2}{3}} \left(\frac{|Q|}{6} \right)^{-\frac{1}{3}} \right] \frac{1}{\ell_p^2}, \quad \text{where} \quad m_2 = \frac{1}{r_{\text{UV}}}.$$

The mass spectrum of the $\bar{v}(x)$ field is calculated similarly

$$M_{\bar{v}\bar{v}}^2(N_{\text{II}}^{0,1,0}) \equiv \left(\frac{\partial^2}{\partial \bar{v}^2} 2\bar{V}_2 \right)_{\bar{u}=\bar{u}_2, \bar{v}=\bar{v}_2} = \frac{20}{27} \bar{\Lambda}_2 \ell_p^2 = -\frac{20}{9} m_2^2. \quad (3.4)$$

From the mass formula for the $SU(3) \times SU(2)$ representation and the eigenvalues of the Lichnerowicz operator, one should obtain the mass-squared for the singlet as follows: $M_{(\bar{\mathbf{1}}, \mathbf{1})}^2 = -\frac{20}{9} m_2^2$, and this coincides with (3.4). The perturbation that corresponds to squashing around $N_{\text{II}}^{0,1,0}$ has a scaling dimension either $\Delta = 4/3$ or $5/3$ and hence corresponds to a relevant operator.

We gave a nonzero expectation value to a supergravity scalar in the $\bar{\mathbf{6}}$ of $SU(3)$. Using the AdS/CFT correspondence, one identifies this perturbation with a composite operator of $\mathcal{N} = 3$ superconformal Chern-Simons matter theory with a mass term for the symmetric and traceless product between two $\bar{\mathbf{3}}$'s: $\lambda^{AB} \int d^3x \mathcal{O}_{AB}$ where λ^{AB} is in the $\mathbf{6}$ of $SU(3)$. Note that the tensor product of these leads to $\bar{\mathbf{3}}(\boxed{}) \times \bar{\mathbf{3}}(\boxed{}) = \mathbf{3}(\square) \oplus \bar{\mathbf{6}}(\boxed{} \oplus \boxed{})$. Then one can construct a $\bar{\mathbf{3}}$ (that is the Young tableaux $\boxed{}$) representation by using the Clebsch-Gordan coefficient Γ_{AIJ} ($A = 1, 2, 3$) which transforms two $\mathbf{3}$'s into $\bar{\mathbf{3}}$ of $SU(3)$ ($\mathbf{3}(\square) \times \mathbf{3}(\square) = \bar{\mathbf{3}}(\boxed{}) \oplus \mathbf{6}(\square \oplus \square)$) together with matter field C^I : $\Gamma_{AIJ} C^I C^J$ where C^I ($I = 1, 2, 3$) are three complex scalars ($\mathbf{3}$ under the $SU(3)$) transforming as $(\mathbf{N}, \bar{\mathbf{N}})$ with gauge group $SU(N) \times SU(N)$ in $\mathcal{N} = 3$ superconformal Chern-Simons gauge theory [19]. The perturbation is given by

$$\mathcal{O}_{AB} \sim \text{Tr} \Gamma_{AIJ} C^I C^J \Gamma_{BKL} C^K C^L.$$

The singlet of this operator $\mathcal{O}_{AB}$ which is $\bar{\mathbf{6}}$ of $SU(3)$ corresponds to the supergravity field $v(x)$ (or $\bar{v}(x)$) and the conformal dimensions are given by $\Delta_{UV} = \frac{4}{3}$ (or $\frac{5}{3}$) and $\Delta_{IR} = 4$ respectively as we computed before. The other five states with $J = 2$ among $\bar{\mathbf{6}}$ of $SU(3)$ are non-diagonal and correspond to deformations of the seven-dimensional metric [23]. Since the Lichnerowicz operator provides nine-dimensional space, the remaining three states correspond to the eigenvalue $M_{\mathbf{200}} = 0$. These are organized in a triplet ($J = 1$) of $SU(2)_R$ and the massless scalars belong to the additional massless vector multiplet. So far we considered only FR compactification where there are no internal components for the four-form field strength. In [23], they further described the deformation from turning on an internal three-form.

4 Conclusions and outlook

We have constructed the full eleven-dimensional metric given by (2.1) and obtained the scalar potential in (2.10) by using the Freund-Rubin ansatz (2.3). The holographic supersymmetric (or non-supersymmetric) RG flow from $\mathcal{N} = (0, 1)$ $SU(3) \times SU(2)$ -invariant UV fixed point

to $\mathcal{N} = (3, 0)$ $SU(3) \times SU(2)$ -invariant IR fixed point was described. The corresponding operator in three-dimensional Chern-Simons matter theories is identified.

For the seven-dimensional Einstein metric, we considered $SO(3)$ -bundle over the base $\mathbf{CP}^2$ in (2.1). Also the base $\mathbf{CP}^2$ can be replaced by either $\mathbf{S}^4$ or $\mathbf{CP}^1 \times \mathbf{CP}^1$. It would be interesting to find out whether these metrics provide the new nontrivial eleven-dimensional solutions. Are there any new general AdS_4 vacua corresponding to any supersymmetric Chern-Simons matter theories? Are there any new critical points in the context of $SO(3)$ gauged supergravity (which might be related to $\mathcal{N} = 3$ Chern-Simons matter theory) or $SO(4)$ gauged supergravity?

Acknowledgments

This work was supported by grant No. R01-2006-000-10965-0 from the Basic Research Program of the Korea Science & Engineering Foundation.

References

- [1] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,” arXiv:0806.1218 [hep-th].
- [2] B. E. W. Nilsson and C. N. Pope, “Hopf Fibration Of Eleven-Dimensional Supergravity,” *Class. Quant. Grav.* **1**, 499 (1984).
- [3] P. G. O. Freund and M. A. Rubin, “Dynamics Of Dimensional Reduction,” *Phys. Lett. B* **97**, 233 (1980).
- [4] G. Jensen, “Einstein metrics on principal fibre bundles,” *J. Diff. Geom.* **8** (1973) 599.
- [5] J. P. Bourguignon and H. Karcher, *Ann. Sci. Normale Sup.* **11** (1978) 71.
- [6] M. A. Awada, M. J. Duff and C. N. Pope, “N = 8 supergravity breaks down to N = 1,” *Phys. Rev. Lett.* **50**, 294 (1983).
- [7] M. J. Duff, B. E. W. Nilsson and C. N. Pope, “Spontaneous Supersymmetry Breaking By The Squashed Seven Sphere,” *Phys. Rev. Lett.* **50**, 2043 (1983); **51**, 846 (1983) (errata).
- [8] L. Girardello, M. Petrini, M. Porrati and A. Zaffaroni, “Novel local CFT and exact results on perturbations of N = 4 super Yang-Mills from AdS dynamics,” *JHEP* **9812**, 022 (1998).

- [9] D. Z. Freedman, S. S. Gubser, K. Pilch and N. P. Warner, *Adv. Theor. Math. Phys.* **3**, 363 (1999).
- [10] C. Ahn and S. J. Rey, “Three-dimensional CFTs and RG flow from squashing M2-brane horizon,” *Nucl. Phys. B* **565**, 210 (2000).
- [11] C. Ahn, “Holographic Supergravity Dual to Three Dimensional N=2 Gauge Theory,” *JHEP* **0808**, 083 (2008).
- [12] C. Ahn, “Towards Holographic Gravity Dual of N=1 Superconformal Chern-Simons Gauge Theory,” *JHEP* **0807**, 101 (2008).
- [13] C. Ahn, “Squashing Gravity Dual of N=6 Superconformal Chern-Simons Gauge Theory,” *arXiv:0809.3684 [hep-th]*.
- [14] H. Ooguri and C. S. Park, “Superconformal Chern-Simons Theories and the Squashed Seven Sphere,” *arXiv:0808.0500 [hep-th]*.
- [15] L. Castellani and L. J. Romans, “N=3 And N=1 Supersymmetry In A New Class Of Solutions For D = 11 Supergravity,” *Nucl. Phys. B* **238**, 683 (1984).
- [16] L. Castellani, “The Mass Spectrum In The SU(3) X U(1) Compactifications Of D = 11 Supergravity,” *Nucl. Phys. B* **254**, 266 (1985).
- [17] D. N. Page and C. N. Pope, “New Squashed Solutions Of D = 11 Supergravity,” *Phys. Lett. B* **147**, 55 (1984).
- [18] C. Ahn and S. J. Rey, “More CFTs and RG flows from deforming M2/M5-brane horizon,” *Nucl. Phys. B* **572**, 188 (2000).
- [19] M. Billo, D. Fabbri, P. Fre, P. Merlatti and A. Zaffaroni, “Rings of short N = 3 superfields in three dimensions and M-theory on AdS(4) x N(0,1,0),” *Class. Quant. Grav.* **18**, 1269 (2001).
- [20] D. Gaiotto and X. Yin, “Notes on superconformal Chern-Simons-matter theories,” *JHEP* **0708**, 056 (2007).
- [21] P. Termonia, “The complete N = 3 Kaluza-Klein spectrum of 11D supergravity on AdS(4) x N(010),” *Nucl. Phys. B* **577**, 341 (2000).

- [22] P. Fre', L. Gualtieri and P. Termonia, "The structure of $N = 3$ multiplets in $AdS(4)$ and the complete $Osp(3-4) \times SU(3)$ spectrum of M-theory on $AdS(4) \times N(0,1,0)$," *Phys. Lett. B* **471**, 27 (1999).
- [23] M. Billo, D. Fabbri, P. Fre, P. Merlatti and A. Zaffaroni, "Shadow multiplets in $AdS(4)/CFT(3)$ and the super-Higgs mechanism," *Nucl. Phys. B* **591**, 139 (2000).
- [24] D. Fabbri and P. Fre, "Shadow multiplets and superHiggs mechanism," *Fortsch. Phys.* **49**, 475 (2001).
- [25] B. S. Acharya, J. M. Figueroa-O'Farrill, C. M. Hull and B. J. Spence, "Branes at conical singularities and holography," *Adv. Theor. Math. Phys.* **2**, 1249 (1999).
- [26] D. R. Morrison and M. R. Plesser, "Non-spherical horizons. I," *Adv. Theor. Math. Phys.* **3**, 1 (1999).
- [27] D. L. Jafferis and A. Tomasiello, "A simple class of $N=3$ gauge/gravity duals," arXiv:0808.0864 [hep-th].
- [28] K. M. Lee and H. U. Yee, "New $AdS(4) \times X(7)$ geometries with $N = 6$ in M theory," *JHEP* **0703**, 012 (2007).
- [29] B. M. Zupnik and D. V. Khetselius, "Three-dimensional extended supersymmetry in harmonic superspace," *Sov. J. Nucl. Phys.* **47** (1988) 730 [*Yad. Fiz.* **47** (1988) 1147].
- [30] H. C. Kao, "Selfdual Yang-Mills Chern-Simons Higgs systems with an $N=3$ extended supersymmetry," *Phys. Rev. D* **50**, 2881 (1994).
- [31] H. C. Kao, K. M. Lee and T. Lee, "The Chern-Simons coefficient in supersymmetric Yang-Mills Chern-Simons theories," *Phys. Lett. B* **373**, 94 (1996).
- [32] M. J. Duff, B. E. W. Nilsson and C. N. Pope, "Kaluza-Klein Supergravity," *Phys. Rept.* **130**, 1 (1986).
- [33] C. Ahn, "More Penrose limit of $AdS(4) \times N(0,1,0)$ and $N = 3$ gauge theory," *Mod. Phys. Lett. A* **17**, 1847 (2002).
- [34] D. N. Page, "Classical Stability Of Round And Squashed Seven Spheres In Eleven-Dimensional Supergravity," *Phys. Rev. D* **28**, 2976 (1983).
- [35] V. L. Campos, G. Ferretti, H. Larsson, D. Martelli and B. E. W. Nilsson, "A study of holographic renormalization group flows in $d = 6$ and $d = 3$," *JHEP* **0006**, 023 (2000).

- [36] O. Yasuda, “Classical Stability Of M (Pqr), Q (Pqr) And N (Pqr) In D = 11 Supergravity,” Phys. Rev. Lett. **53**, 1207 (1984).
- [37] J. P. Gauntlett, S. Lee, T. Mateos and D. Waldram, “Marginal deformations of field theories with AdS(4) duals,” JHEP **0508**, 030 (2005).
- [38] R. Slansky, “Group Theory For Unified Model Building,” Phys. Rept. **79**, 1 (1981).
- [39] J. Patera and D. Sankoff, *Tables of Branching Rules for Representations of Simple Lie Algebras*(L’Université de Montréal, Montréal, 1973); W. MacKay and J. Petera, *Tables of Dimensions, Indices and Branching Rules for representations of Simple Algebras*(Dekker, New York, 1981).
- [40] R. D’Auria and P. Fre, “Universal Bose-Fermi Mass Relations In Kaluza-Klein Supergravity And Harmonic Analysis On Coset Manifolds With Killing Spinors,” Annals Phys. **162**, 372 (1985).
- [41] I. Klebanov, T. Klose and A. Murugan, “ AdS_4/CFT_3 – Squashed, Stretched and Warped,” arXiv:0809.3773 [hep-th].